Supplementary Table 1. ICIQ-FLUTS scores according to daily screen exposure duration in female participants

Daily screen exposure duration	n	ICIQ-FLUTS score, median (IQR)
<1 hour	4	7.00 (4.50-11.50)
1-2 hours	26	6.00 (2.00-8.00)
2-4 hours	80	5.00 (3.00-7.00)
>4 hours	187	5.00 (3.00-9.00)

Data are presented as median (IQR). P-value was calculated using the Kruskal-Wallis test (p=0.304)
IQR: Interquartile range, ICIQ-FLUTS: International Consultation on Incontinence Questionnaire female lower urinary tract symptoms

Supplementary Table 2. ICIQ-MLUTS scores according to daily screen exposure duration in male participants

Daily screen exposure duration	n	ICIQ-MLUTS score, median (IQR)
<1 hour	1	8.00 (8.00-8.00)
1-2 hours	14	5.00 (3.00-7.00)
2-4 hours	56	4.00 (2.00-7.00)
>4 hours	110	5.00 (3.00-8.00)

Data are presented as median (IQR). P-value was calculated using the Kruskal-Wallis test (p=0.476)
IQR: Interquartile range, ICIQ-MLUTS: International Consultation on Incontinence Questionnaire male lower urinary tract symptoms