

RAD51, BRCA1, and BRCA2: Can be Promising Prognostic Factors and New Therapy Targets for Clear Cell Renal Cell Carcinoma Patients?

Amira M. Abd El Maged¹, Heba Mahmoud Abdelgeleel², Eslam G. Abozaid³, Amira H. Hanna⁴, Hanan Lotfy Mohammed⁵

¹Menoufia University Faculty of Medicine, Shebin El Kom, Egypt

²Zagazig University Faculty of Medicine, Department of Pathology, Zagazig, Egypt

³Zagazig University Surgical Oncology Unit, Department of General Surgery, Zagazig, Egypt

⁴Zagazig University Faculty of Medicine, Department of Clinical Oncology and Nuclear Medicine, Zagazig, Egypt

⁵Zagazig University Faculty of Medicine, Zagazig, Egypt and Ibn Sina National College for Medical Studies (ISNC), Department of Pathology, Jeddah, Saudi Arabia

What's known on the subject? and What does the study add?

RAD51 showed increased immunohistochemical (IHC) expression in clear cell renal cell carcinoma (ccRCC), while *BRCA1* and *BRCA2* show reduced IHC expression, suggesting impaired homologous recombination (HR) repair mechanisms in ccRCCs. High *RAD51* expression is associated with poor prognostic factors of ccRCC patients. High *BRCA1/BRCA2* expression is linked to good prognostic factors of ccRCC patients. Expression of these DNA repair proteins may serve as prognostic markers for ccRCC patients, helping to predict cancer aggressiveness and metastasis and detect outcomes. Targeting HR-proficient /deficient ccRCC with DNA repair-targeting therapies, *RAD51* inhibitors or PARP inhibitors could be a promising treatment approach.

Abstract

Objective: Clear cell renal cell carcinoma (ccRCC) is the predominant form of renal cell carcinoma, distinguished by its genetic alterations that affects the tumor's progression and response to treatment. *RAD51*, *BRCA1*, and *BRCA2* are fundamental components of the homologous recombination (HR) DNA repair system, which is vital for preserving genomic stability. Yet, data regarding their expression pattern and potential role in ccRCC development and progression remain unclear. The aim of this work was to study the immunohistochemical expression of *RAD51*, *BRCA1*, and *BRCA2* in ccRCC and to investigate the correlation between their expression levels and the clinical outcomes of patients with ccRCC.

Materials and Methods: Across-sectional study has been performed at Zagazig University's Faculty of Medicine using data from the Pathology Department archives between January 2020 and December 2023 through analyzing 76 ccRCC specimens who underwent either a total (56 cases) or partial nephrectomy (20 cases) to determine the *RAD51*, *BRCA1*, and *BRCA2* immunohistochemical expressions and comparing these with clinicopathologic parameters and survival.

Results: High nuclear expression of *RAD51* was detected in 42.1% of cases and was significantly associated with poor prognostic indicators, including advanced tumor stage, lymphatic spread, distant metastasis, and recurrence. High nuclear expression of *BRCA1*, *BRCA2*, and both proteins was detected in 51.3%, 35.5%, and 27% of cases, respectively, and their immune expression was significantly correlated with favorable prognostic indicators, including early tumor stage and absence of lymphatic spread. High *RAD51* expression showed significant indirect correlations with *BRCA1* expression and with the co-expression of *BRCA1* and *BRCA2*. Using Kaplan-Meier analysis, patients with high *RAD51* and low *BRCA1* expression had shorter disease-free survival and overall survival.

Conclusion: This research emphasizes the significance of *RAD51*, *BRCA1*, and *BRCA2* in the progression of ccRCC. Elevated *RAD51* levels alongside reduced *BRCA1* and *BRCA2* expression are associated with more aggressive tumors, suggesting their potential as immunohistochemical prognostic indicators and providing insights into novel therapeutic approaches for patients with ccRCC.

Keywords: *RAD51*, *BRCA1*, *BRCA2*, clear cell renal cell carcinoma, immunohistochemistry, homologous recombination, prognosis

Correspondence: Heba Mahmoud Abdelgeleel MD, Zagazig University Faculty of Medicine, Department of Pathology, Zagazig, Egypt

E-mail: HMAbduljalil@medicine.zu.edu.eg **ORCID-ID:** orcid.org/0000-0002-2508-1770

Received: 08.09.2025 **Accepted:** 04.11.2025 **Epub:** 13.07.2026 **Publication Date:** 01.09.2026

Cite this article as: Abd El Maged AM, Abdelgeleel HM, Abozaid EG, Hanna AH, Mohammed HL. *RAD51, BRCA1, and BRCA2: can be promising prognostic factors and new therapy targets for clear cell renal cell carcinoma patients?* J Urol Surg. 2026;13(3):186-198

©Copyright 2026 The Author(s). Published by Galenos Publishing House on behalf of the Society of Urological Surgery.

This is an open access article under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND) International License.



Introduction

Renal cell carcinoma (RCC) is a highly prevalent malignancy of the urinary system and one of the ten most frequently diagnosed solid tumors (1). Clear cell renal cell carcinoma (ccRCC) is the predominant histopathological subtype, constituting more than 80% of RCCs (2). While ccRCC generally has a good prognosis when treated in its early stages, up to one-third of cases will metastasize, and 20-40% of cases will experience a post-nephrectomy recurrence. ccRCC is notably resistant to chemotherapy, and once recurrence occurs, no further management is effective (3).

Deletion of chromosome 3p and mutations in the *Von Hippel-Lindau (VHL)* gene are prevalent in most ccRCC patients, leading to loss of various tumor suppressor genes, which further enhances genomic instability (4). Moreover, a deficiency in homologous recombination (HR) significantly contributes to genomic instability (5).

HR is recognized as a precise process for the repair of DNA double-strand breaks (DSBs), and any defect in this process can lead to tumorigenesis by enabling the accumulation of chromosomal abnormalities and mutations. In numerous instances, aggressive tumors are characterized by reduced expression of HR proteins, but these tumors also show increased responsiveness to therapeutic interventions, including chemotherapy and radiotherapy. The HR pathway involves important proteins encoded by the *RAD51*, *BRCA1*, and *BRCA2* genes (6).

The *RAD51* gene on human chromosome 15q15.1 encodes the essential HR protein *RAD51*. It has a significant role in DSBs by creating nucleoprotein filaments on single-stranded DNA, enabling strand exchange between single- and double-stranded DNA, and facilitating homologous pairing. When DNA damage occurs, *RAD51* relocates to nuclear foci, where it is believed to recognize and facilitate the repair of broken replication forks (7). Recent research has demonstrated that *RAD51* is a potential prognostic biomarker for various tumors, such as colorectal cancer (8), non-small cell lung cancer (9), and breast cancer (10). Nonetheless, there is a lack of evidence demonstrating the link between *RAD51* and ccRCC, which was the rationale for selecting this marker to be investigated in the current study.

The tumor suppressor gene, *BRCA1*, plays a crucial role in maintaining genome stability. It is frequently impaired in several types of cancer, including breast, ovarian, and pancreatic cancers, leading to the accumulation of genetic defects (11). *BRCA1* forms a complex with *BARD1* to aid DSB repair through HR. The complex formed by *BRCA1-BARD1* interacts directly with *RAD51*, boosting its ability to facilitate recombination. Mutations in *BRCA1* that disrupted the *RAD51* interaction hinder DNA strand invasion and compromise HR (12). Therefore, *BRCA1* serves as a crucial prognostic and therapeutic target for advancing cancer treatment approaches.

BRCA2 protein promotes HR-mediated repair of DSBs by facilitating *RAD51*-dependent DNA strand invasion, pairing, and exchange. The C-terminal recombinase binding (CTRB) domain of *BRCA2* is responsible for loading *RAD51* onto and binding *RAD51* to single-stranded DNA (ssDNA). CTRB alone significantly increases the DNA strand-exchange activity of *RAD51* and enables it to use ssDNA for this activity (13). In human cells, *BRCA2* deficiency leads to HR abnormality, genetic instability, inhibited formation of *RAD51* foci in response to DNA damage, and altered responsiveness to chemotherapy agents (14).

This study is the first to analyze immunohistochemical (IHC) expression of *RAD51*, *BRCA1*, and *BRCA2* in ccRCC and to investigate correlations between their expression levels and the clinicopathological features and clinical outcomes in patients with ccRCC. This research may help identify additional prognostic indicators for ccRCC and inform the development of novel anticancer strategies.

Materials and Methods

This cross-sectional study was conducted at Zagazig University's Faculty of Medicine using data from the Pathology Department archives collected from January 2020 to December 2023. The study included 76 sections from formalin-settled paraffin-inserted tissue blocks diagnosed as primary ccRCC who underwent either a total (56 cases) or partial nephrectomy (20 cases). Patients were selected consecutively. Clinicopathologic information, histopathology reports, and hematoxylin and eosin-stained slides from ccRCC cases were reviewed to confirm the diagnosis. Exclusion criteria were: a history of synchronous or metachronous malignancies; prior antitumor therapy; histological types other than the clear-cell variant; or insufficient tissue for IHC analysis. This work was conducted in agreement with the Helsinki Declaration and written consent was obtained from each member. Moreover, the current study was permitted by the Ethical Committee of Zagazig University according to the Egyptian Ethical Guidelines (approval number: ZU-IRB#11335, date: 12.03.2024).

Immunohistochemistry

The immunostaining method employed a streptavidin-biotin amplification system. The slides were subjected to consecutive steps of deparaffinization, rehydration, and blocking of endogenous peroxidase activity. Antigen recovery was performed by boiling in citrate-buffered saline (pH 6), followed by cooling at room temperature. The slides were then stained with primary antibodies against *RAD51* (rabbit monoclonal Ab; Catalog # EPR4030(3); 1:300 dilution; Abcam, Cambridge, UK), *BRCA1* (rabbit monoclonal Ab; Catalog # EPR19433; 1:400 dilution; Abcam, Cambridge, UK), and *BRCA2* (rabbit polyclonal Ab; Catalog # ab216972; 1:400 dilution; Abcam, Cambridge, UK). The

primary antibody was incubated overnight at room temperature, and then the secondary antibody was applied; diaminobenzidine was used as the chromogen, followed by counterstaining with Mayer's hematoxylin. The *RAD51* positive-control slide used germ cells from healthy testicular tissue. The positive control for *BRCA1* and *BRCA2* was tissue from invasive ductal carcinoma of the breast. Negative control slides were treated without the addition of primary antibodies.

Immunohistochemical Evaluation Criteria

Two independent investigators examined all stained sections by light microscopy; interobserver agreement was achieved. They specifically focused on assessing the nuclear expression of *RAD51*, *BRCA1*, and *BRCA2*. During the pathological annotation process, the assessment involved determining the percentage of positive cells, categorized as follows: 0 for less than 5%, 1 for 5-25%, 2 for 25-50%, 3 for 50-75%, and 4 for 75-100%. Additionally, the intensity of staining was evaluated on a scale where 0 indicates negative, 1 represents weak, 2 denotes moderate, and 3 signifies strong staining. The overall score was derived by multiplying the intensity score by the percentage score. Samples were subsequently classified as either "high expression" or "low expression" based on a cut-off score of 6 (15).

Analysis of Survival Data

For 76 patients with ccRCC between January 2020 and December 2023, the overall survival (OS) time was determined by reviewing the patients' files. Disease-free survival (DFS) and OS were analyzed using the Kaplan-Meier method, and differences among the survival curves were assessed by a log-rank test. A multivariate Cox proportional hazards analysis was employed to determine the independent risk factors influencing survival time.

Statistical Analysis

SPSS 22.0 for Windows and MedCalc Windows were used to perform all analyses. The Mann-Whitney U test was employed to compare two groups of non-normally distributed variables, whereas the Kruskal-Wallis test was utilized for comparisons involving more than two groups of non-normally distributed variables. The proportions of categorical variables were compared using Fisher's exact test or Pearson's chi-square test, as appropriate. A p-value ≤ 0.05 was considered statistically significant.

Results

Patients' Characteristics

This study included 76 patients with ccRCC. Patients' ages ranged from 48-80 years, with a mean age of 58.66 ± 7.561 years (Table 1). There were 45 (59.2%) male patients and 31 (40.8%) female patients. Tumor size exceeded 7 cm in 54 cases (71.1%).

Thirty-nine (51.3%) cases were high grade, while 37 (48.7%) were low grade. Necrosis was detected in 31 (40.8%) cases, while lymphovascular invasion was present in 19 (25%) cases. Seventeen of these 76 patients (22.4%) had stage I disease. On the other hand, 20 ccRCC patients (26.3%) presented with stage IV disease. Nodal metastasis was recorded preoperatively in 25 (32.9%) and distant metastasis in 23 (30.3%) cases as well. The average duration of follow-up was 27.4 months (range: 10-36 months), during which 40 patients (52.6%) remained disease-free. Recurrence occurred in 36 patients (47.4%), while 28 patients (36.8%) died during the follow-up period. The clinicopathological features of the 76 ccRCC patients are detailed in Table 1.

Pattern of *RAD51* Expression in ccRCC and Association with Clinicopathological Parameters

In ccRCC patients, nuclear staining of tumor cells was considered *RAD51*-positive (Figure 1). High *RAD51* expression was observed in 32 patients (42.1%), while 44 patients (57.9%) showed low expression (Table 1). In adjacent non-tumor tissue, no immunostaining for *RAD51* was observed.

Table 1. Clinicopathologic parameters of 76 ccRCC cases	
Variable	No (%)
Gender	
Male	45 (59.2)
Female	31 (40.8)
Age	
Mean $\pm$ standard deviation	58.66 $\pm$ 7.561
Median	57
Range	32
Age	
<50	20 (26.3)
≥ 50	56 (73.7)
Tumor size	
<7	22 (28.9)
≥ 7	54 (71.1)
The WHO/ISUP grade	
Grade I-II	37 (48.7)
Grade III-IV	39 (51.3)
Vascular invasion	
Present	19 (25)
Absent	57 (75)
Necrosis	
Present	31 (40.8)
Absent	45 (59.2)
Tumor stage	
T1	17 (22.4)
T2	23 (30.3)

Table 1. Continued	
Variable	No (%)
T3	16 (21.1)
T4	20 (26.3)
Tumor stage grouping	
Early stage	40 (52.6)
Advanced stage	36 (47.4)
Nodal stage	
N0	51 (67.1)
N1	25 (32.9)
Distant metastasis	
M0	53 (69.7)
M1	23 (30.3)
Relapse	
Absent	40 (52.6)
Present	36 (47.4)
Death	
Alive	48 (63.2)
Dead	28 (36.8)
<i>RAD51</i>	
Low expression	44 (57.9)
High expression	32 (42.1)
<i>BRCA1</i>	
Low expression	37 (48.7)
High expression	39 (51.3)
<i>BRCA2</i>	
Low expression	49 (64.5)
High expression	27 (35.5)
High co-expression of <i>BRCA1</i> and <i>BRCA2</i>	
Positive	21 (27.6)
Negative	55 (72.4)
ISUP: International Society of Urologic Pathologists, WHO: World Health Organization, ccRCC: Clear cell renal cell carcinoma	

High *RAD51* expression was significantly correlated with higher tumor stage, with high expression observed in 4 (23.5%) stage I, 6 (26.1%) stage II, 8 (50%) stage III, and 14 (70%) stage IV ccRCC cases ($p=0.009$); the correlation was significant ($p=0.001$). Moreover, high *RAD51* was significantly associated with tumor lymphatic spread ($p<0.001$ and $p=0.000$), distant spread ($p<0.001$ and $p=0.000$), and tumor recurrence ($p=0.024$ and $p=0.02$) (Table 2, Figure 1).

Pattern of *BRCA1* and *BRCA2* Expression in ccRCC and Association with Clinicopathological Parameters

BRCA1 and *BRCA2* showed nuclear staining in cancer cells, with negative immunostaining in the adjacent non-tumor

tissue (Figure 2). High immunohistochemical (IHC) expression of *BRCA1*, *BRCA2*, and both proteins were observed in 39 (51.3%), 27 (35.5%), and 21 (27.6%) cases, respectively (Table 1).

BRCA1 immune expression was significantly correlated with early stages of ccRCC, with higher expression observed in 14 (82.4%) stage I cases compared with 8 (40%) stage IV cases ($p=0.017$); the correlation was significant ($p=0.006$). Additionally, *BRCA1* expression was significantly associated with cases without lymphatic spread ($p=0.028$). Similarly, co-expression of *BRCA1* and *BRCA2* was significantly associated with early stages of ccRCC patients ($p=0.004$; $p=0.002$), and lack of nodal metastasis ($p=0.032$; $p=0.01$). However, *BRCA2* expression is significantly correlated with the early tumor stage ($p=0.008$) and with the correlation coefficient ($p=0.005$) (Table 3; Figure 2).

Association Between *RAD51*, *BRCA1*, and *BRCA2* Expression

A significant correlation was detected between positive *RAD51* expression and negative *BRCA1* expression: 30 (68.2%) cases with high *BRCA1* expression showed low *RAD51* expression, whereas 9 (28.1%) of the high *BRCA1* expression cases showed high *RAD51* immunostaining ($p=0.001$; correlation coefficient $p=0.000$). No significant correlation was found between *RAD51* and *BRCA2* immune expression ($p=0.507$); however, co-expression of *BRCA1* and *BRCA2* showed a significant inverse association with *RAD51*. In the correlation between *BRCA1* and *BRCA2*, a significant positive association was detected ($p=0.001$); 21 cases (27.6%) with high *BRCA2* expression also showed high *BRCA1* expression, and the correlation coefficient was significant ($p<0.001$) (Figure 3).

Univariate Survival Analysis for Predicting Overall & Disease-free Survival

Univariate analysis of OS and DFS in 76 ccRCC patients using the Kaplan-Meier method (Figure 4) revealed that high tumor grade ($p=0.001$ and 0.001), advanced stage ($p=0.043$ and 0.016), presence of nodal metastasis ($p=0.004$ and 0.003), distant metastasis ($p=0.02$ and 0.008), and tumor recurrence ($p<0.001$ and <0.001) were associated with shorter OS and DFS, respectively (Table 4).

High expression of *RAD51* was associated with shorter OS and DFS ($p=0.02$ and $p=0.014$, respectively). Cases with a mean OS of 30.93 months exhibited high *RAD51* immunostaining, whereas cases with a longer mean OS of 32.04 months displayed lower *RAD51* expression levels. Conversely, *BRCA1* expression was significantly associated with longer OS and DFS ($p=0.04$ and $p=0.009$, respectively). However, *BRCA2* showed no significant correlation with patient survival (Figure 4, Table 5).




Figure 1. I) High and low *RAD51* immunohistochemical expression in clear cell renal cell carcinoma cases (magnified x400). **II)** High *RAD51* showed a significant association with advanced tumor stage (A), nodal involvement (B), distant metastasis (C), and tumor relapse (D)

Table 2. Correlation between <i>RAD51</i> with clinicopathologic parameters of 76 ccRCC cases			
Variable	<i>RAD51</i> expression		p-value
	Low expression No (%)	High expression No (%)	
Gender			0.357
Male	(28) 62.2%	(17) 37.8%	
Female	(16) 51.6%	(15) 48.4%	
Age			0.453
<50	(13) 65.0%	(7) 35.0%	
≥50	(31) 55.4%	(25) 44.6%	
Tumor size			0.706
<7	(12) 54.5%	(10) 45.5%	
≥7	(32) 59.3%	(22) 40.7%	
WHO/ISUP grade			0.231
Grade I-II	(24) 64.9%	(13) 35.1%	
Grade III-IV	(20) 51.3%	(19) 48.7%	
Vascular invasion			0.592
Present	(10) 52.6%	(9) 47.4%	
Absent	(34) 59.6%	(23) 40.4%	
Necrosis			0.164
Present	(15) 48.4%	(16) 51.6%	
Absent	(29) 64.4%	(16) 35.6%	
Tumor stage			0.009*
T1	(13) 76.5%	(4) 23.5%	
T2	(17) 73.9%	(6) 26.1%	
T3	(8) 50%	(8) 50%	
T4	(6) 30%	(14) 70%	
Tumor stage grouping			0.001*
Early stage	(30) 75%	(10) 25%	
Advanced stage	(14) 38.9%	(22) 61.1%	<0.001*
Nodal stage			
N0	(37) 72.5%	(14) 27.5%	
N1	(7) 28%	(18) 72%	

Table 2. Continued			
Variable	RAD51 expression		p-value
	Low expression No (%)	High expression No (%)	
Distant metastasis			<0.001*
M0	(38) 71.7%	(15) 28.3%	
M1	(6) 26.1%	(17) 53.1%	
Relapse			0.024*
Absent	(28) 70%	(12) 30%	
Present	(16) 44.4%	(20) 55.6%	
BRCA1			0.001*
Low expression	(14) 37.8%	(23) 62.2%	
High expression	(30) 76.9%	(9) 23.1%	
BRCA2			0.507
Low expression	(27) 55.1%	(22) 44.9%	
High expression	(17) 63%	(10) 37%	
High co-expression of BRCA1 and BRCA2			0.018*
Positive	(17) 81%	(4) 19%	
Negative	(27) 49.1%	(28) 50.9%	
*: Significant, ISUP: International Society of Urologic Pathologists, WHO: World Health Organization, ccRCC: Clear cell renal cell carcinoma			

*: Significant, ISUP: International Society of Urologic Pathologists, WHO: World Health Organization, ccRCC: Clear cell renal cell carcinoma




Figure 2. I) High and low *BRCA1* and *BRCA2* immunohistochemical expression in clear cell renal cell carcinoma cases (magnified x400). **II)** High *BRCA1* showed significant associations with early tumor stage (A) and with presence of nodal stage (B); *BRCA2* showed significant association with early tumor stage (C). Co-expression of both showed significant association with early tumor stage (D), and presence of nodal stage (E)

Variable	<i>BRCA1</i> expression		p-value	<i>BRCA2</i> expression		p-value	Co-expression		p-value
	Low expression No (%)	High expression No (%)		Low expression No (%)	High expression No (%)		Negative expression No (%)	Positive expression No (%)	
Gender			0.672			0.142			0.07
Male	(21) 46.7%	(24) 53.3%		(26) 57.8%	(19) 42.2%		(29) 64.4%	(16) 35.6%	
Female	(16) 51.6%	(15) 48.4%		(23) 74.2%	(8) 25.8%		(26) 83.9%	(5) 16.1%	
Age			0.701			0.302			0.78
<50	(9) 45.0%	(11) 55.0%		(11) 55%	(9) 45%		(14) 70%	(6) 30%	
≥50	(28) 50%	(28) 50%		(38) 67.9%	(18) 32.1%		(41) 73.2%	(15) 26.8%	
Tumor size			0.387			0.248			0.60
<7	(9) 40.9%	(13) 59.1%		(12) 54.5%	(10) 45.5%		(15) 68.2%	(7) 31.8%	
≥7	(28) 51.9%	(26) 48.1%		(37) 68.5%	(17) 31.5%		(40) 74.1%	(14) 25.9%	
WHO/ISUP grade			0.167			0.171			0.15
Grade I-II	(15) 40.5%	(22) 59.5%		(21) 56.8%	(16) 43.2%		(24) 64.9%	(13) 35.1%	
Grade III-IV	(22) 56.4%	(17) 43.6%		(28) 71.8%	(11) 28.2%		(31) 79.5%	(8) 20.5%	
Vascular invasion			0.691			0.128			0.076
Present	(10) 52.6%	(9) 47.4%		(15) 78.9%	(4) 21.1%		(17) 89.5%	(2) 10.5%	
Absent	(27) 47.4%	(30) 52.6%		(34) 59.6%	(23) 40.4%		(38) 66.7%	(19) 33.3%	
Necrosis			0.373			0.326			0.41
Present	(17) 54.8%	(14) 45.2%		(22) 71%	(9) 29%		(24) 77.4%	(7) 22.6%	
Absent	(20) 44.4%	(25) 55.6%		(27) 60%	(18) 40%		(31) 68.9%	(14) 31.1%	
Tumor stage			0.017*			0.03*			0.021*
T1	(3) 17.6%	(14) 82.4%		(7) 41.2%	(10) 58.8%		(10) 58.8%	(7) 41.2%	
T2	(11) 47.8%	(12) 52.2%		(13) 56.5%	(10) 43.5%		(13) 56.5%	(10) 43.5%	
T3	(11) 68.8%	(5) 31.2%		(13) 81.2%	(3) 18.8%		(15) 93.8%	(1) 6.2%	
T4	(12) 60%	(8) 40%		(16) 80%	(4) 20%		(17) 85%	(3) 15%	
Tumor stage grouping			0.021*			0.008*			0.004*
Early	(14) 35%	(26) 65%		(20) 50%	(20) 50%		(23) 57.5%	(17) 42.5%	
Advanced	(23) 63.9%	(13) 36.1%		(29) 80.6%	(7) 19.4%		(32) 88.9%	(4) 11.1%	
Nodal stage			0.028*			0.07			0.032*
N0	(20) 39.2%	(31) 60.8%		(29) 56.9%	(22) 43.1%		(33) 64.7%	(18) 35.3%	
N1	(17) 68%	(8) 32%		(20) 80%	(5) 20%		(22) 88%	(3) 12%	
Distal metastasis			0.080			0.098			0.09
M0	(22) 41.5%	(31) 58.5%		(27) 50.0%	(27) 50.0%		(35) 66%	(18) 34%	
M1	(15) 65.2%	(8) 34.8%		(10) 45.5%	(12) 54.5%		(20) 87%	(3) 13%	
Relapse			0.498			0.561			0.58
Absent	(18) 45%	(22) 55%		(27) 67.5%	(13) 32.5%		(30) 75%	(10) 25%	
Present	(19) 52.8%	(17) 47.2%		(22) 61.1%	(14) 38.9%		(25) 69.4%	(11) 30.6%	
<i>RAD51</i>			0.001*			0.507			0.018
Low expression	(14) 31.8%	(30) 68.2%		(27) 61.4%	(17) 38.6%		(27) 61.4%	(17) 38.6%	
High expression	(23) 71.9%	(9) 28.1%		(22) 68.8%	(10) 31.2%		(28) 87.5%	(4) 12.5%	
ISUP: International Society of Urologic Pathologists, WHO: World Health Organization, ccRCC: Clear cell renal cell carcinoma									

ISUP: International Society of Urologic Pathologists, WHO: World Health Organization, ccRCC: Clear cell renal cell carcinoma




Figure 3. *RAD51* expression showed significantly inverse association with *BRCA1* (A), and co-expression of *BRCA1* & *BRCA2* (B). Direct significant association between *BRCA1* and *BRCA2* (C)




Figure 4. Kaplan-Meier plot, upper panel: Overall survival, lower panel: Disease free survival. (A&F) all studied patient: (A&D) stratified by *RAD51*, (B&E) stratified by *BRCA1*, and (C&F) stratified by *BRCA2*

Table 4. Univariate survival analysis for predicting overall & disease-free survival						
Variable	OS			DFS		
	Mean (CI 95%)	Log-rank	p-value	Mean (CI 95%)	Log-rank	p-value
Tumor size						
<7	30.24 (26.09-34.39)	0.132	0.716	27.92 (22.23-33.61)	0.002	0.965
≥7	31.81 (29.79-33.83)			27.80 (24.76-30.84)		
WHO/ISUP grade						
Grade I-II	33.77 (31.58-35.96)	10.78	0.001*	32.21 (29.17-35.25)	11.46	0.001*
Grade III-IV	29.58 (26.87-32.29)			24.59 (20.87-28.30)		
Vascular invasion						
Present	29.32 (24.57-34.06)	0.668	0.414	26.72 (20.25-33.18)	0.44	0.504
Absent	32.17 (30.34-34.00)			28.28 (25.38-31.17)		
Necrosis						
Present	29.80 (26.76-32.85)	2.8	0.094	24.45 (20.08-28.82)	3.45	0.063
Absent	32.74 (30.58-34.91)			30.39 (27.21-33.57)		
Tumor stage						
T1	31.84 (27.68-36.01)	8.14	0.043*	29.74 (23.46-36.03)	10.38	0.016*
T2	33.75 (31.04-36.46)			32.20 (28.42-35.98)		
T3	31.2 (27.49-34.91)			27.46 (21.66-33.29)		
T4	28.92 (24.87-32.98)			22.36 (17.25-27.47)		
Tumor stage grouping						
Early stage	33.10 (30.80-35.40)	7.172	0.007*	31.37 (28.05-34.70)	8.77	0.003*
Advanced stage	29.96 (27.22-32.70)			24.74 (20.91-28.57)		
Nodal stage						
N0	32.71 (30.69-34.73)	8.16	0.004*	30.61 (27.64-33.57)	9.06	0.003*
N1	29.23 (25.79-32.68)			23.15 (18.62-27.67)		
Distal metastasis						
M0	32.48 (30.47-34.49)	5.35	0.02*	30.28 (27.33-33.277)	7.01	0.008*
M1	29.42 (25.79-33.05)			22.97 (18.17-27.78)		
Relapse						
Absent	33.45 (31.26-35.64)	16.05	<0.001*	32.10 (29.25-34.95)	17.26	<0.001*
Present	29.30 (26.44-32.16)			23.28 (19.31-27.25)		
RAD51						
Low expression	32.04 (29.49-34.60)	5.27	0.02*	29.96 (26.40-33.52)	6.02	0.014*
High expression	30.93 (28.41-33.46)			25.57 (21.67-29.46)		
BRCA1						
Low expression	30.52 (27.91-33.12)	3.93	0.04*	25.30 (21.56-29.05)	6.86	0.009*
High expression	32.40 (29.85-34.96)			30.79 (27.23-34.35)		
BRCA2						
Low expression	30.66 (28.25-33.07)	0.72	0.39	26.71 (23.35-30.08)	0.65	0.41
High expression	32.79 (30.12-35.45)			30.03 (25.66-34.41)		
High co-expression of BRCA1 and BRCA2						
Positive	33.56 (30.69-36.44)	2.70	0.10	32.00 (27.49-36.50)	3.46	0.06
Negative	30.67 (28.46-32.93)			26.399 (23.20-29.59)		
OS: Overall survival, DFS: Disease free survival, CI: Confidence interval, *: Significant, ISUP: International Society of Urologic Pathologists, WHO: World Health Organization, ccRCC: Clear cell renal cell carcinoma						

Table 5. Multivariate COX-regression analysis for detection of the most independent prognostic factor affecting patient overall survival		
Variable	Multivariate	
	HR (LL-UL CI 95%)	p-value
WHO/ISUP grade	0.38 (0.104-1.44)	0.158
Tumor stage	0.069 (0.009-0.52)	0.82
Nodal stage	1.13 (0.37-3.39)	0.01*
Distal metastasis	13.12 (1.96-87.68)	0.008*
Relapse	1.32 (0.52-3.33)	0.058
<i>RAD51</i>	0.78 (0.32-1.90)	0.59
<i>BRCA1</i>	0.30 (0.09-1.03)	0.56

HR: Homologous recombination, CI: Confidence interval, LL: Lower limit, UL: Upper limit, *: Significant, ISUP: International Society of Urologic Pathologists, WHO: World Health Organization

Multivariate COX-regression Analysis for Detection of the Most Independent Prognostic Factor Affecting Patient Overall Survival

Nodal metastasis ($p=0.01$) and distant metastasis ($p=0.008$) emerged as the most significant independent prognostic factors affecting OS in ccRCC patients in multivariate Cox proportional-hazards analysis.

Discussion

Genomic instability contributes to tumor initiation and advancement, often resulting from defects in DNA damage repair (DDR) (16). HR, a key DDR pathway, maintains genome stability by repairing DNA DSBs. Due to its role in maintaining genome stability, HR is generally considered a pathway that suppresses tumors. In line with this concept, mutations or deficiencies that result in loss of HR function can also lead to genomic instability and tumor initiation (17).

RAD51, *BRCA1*, and *BRCA2* are crucial proteins in the HR pathway (6), and their relevance to ccRCC has received growing attention as understanding of ccRCC genetics and molecular mechanisms advances. Additionally, their function in DNA repair and cancer development is becoming clearer.

RAD51, a crucial participant in HR, is directly responsible for preserving the stability of the genome. *RAD51* encircles a single broken DNA strand with the assistance of auxiliary proteins, then captures the complementary copy of DNA, aligning it with the sequence of the broken strand (18). Since *RAD51* is essential for maintaining genome stability, *RAD51* expression must be tightly regulated in human cells. Inadequate *RAD51* expression could result in accumulation of DNA breaks. Conversely, an increase in *RAD51* expression could stimulate hyper-recombination, which could cause genomic instability (19).

Studies have reported conflicting findings regarding the prognostic significance of *RAD51* across different cancer types.

In breast cancer (20) and non-small cell lung cancer (21), low *RAD51* expression has been correlated with an unfavorable prognosis. Conversely, high *RAD51* expression has been linked to a poor prognosis in colon cancer (8), breast cancer (10), and non-small-cell lung cancer (9).

This study focused on *RAD51* IHC expression in ccRCC and revealed *RAD51* overexpression in tumor cells. High nuclear *RAD51* expression was observed in a considerable proportion (42.1%) of cases and was significantly associated with advanced tumor stages, nodal metastasis, distant metastasis, and disease recurrence. Survival analysis revealed an association between high *RAD51* expression and shorter OS and DFS in patients with ccRCC. These findings support the tumor-promoting role of *RAD51* and its potential as a novel predictive biomarker in ccRCC patients.

Consistent with our results, *RAD51* overexpression has been consistently observed in multiple cancer types across several studies and has been linked to aggressive proliferation and increased metastatic potential. In a study involving 70 neuroblastoma cases, *RAD51* expression was notably elevated in stage IV tumors compared with stages I and II, particularly in cases with bone marrow metastases (22). A previous study reported an association between high expression of *RAD51* and breast cancer with lymph node metastases (23). Furthermore, Qiao et al. (9) established *RAD51* expression as a potential prognostic indicator in lung cancer, showing that high *RAD51* expression in tumors predicts poor patient survival.

Lu et al. (18) analyzed *RAD51* expression in various cancers, including ccRCC. They found that the expression of *RAD51* was notably elevated in cancerous cells compared with adjacent tissues. Furthermore, the research indicated a strong association between *RAD51* expression and both advanced pathological stages and DFS. Moreover, the study suggested that the expression of *RAD51* was linked to several important immune checkpoints and immunosuppressive genes, indicating its potential impact on controlling the immune response in tumors through regulation of immune checkpoint activity.

RAD51, despite its vital function in the repair of damaged DNA and its possible role as a tumor suppressor, has been linked to genomic instability and the onset of cancer when overexpressed. Increased *RAD51* expression can lead to accumulation of genotoxic *RAD51* on undamaged chromatin, resulting in reduced HR efficiency (24). Additionally, *RAD51* overexpression can positively regulate cell proliferation, decrease intracellular reactive oxygen species production, and regulate aerobic glycolysis by targeting hypoxia-inducible factor 1 α , ultimately contributing to tumor progression (25). *RAD51* is also known to increase the activity of transcription factors, thereby promoting epithelial-mesenchymal transition and production of metalloproteinases (26).

Overexpression of *RAD51* in malignant tissue is linked to a marked rise in HR activity, enhancing tumor cell genetic flexibility. Furthermore, it is believed that it improves the ability to repair DNA breaks, thereby protecting cells from DNA-damaging treatments and conferring resistance to conventional cancer therapies. Consequently, inhibiting *RAD51* to render HR-proficient tumor cells HR-deficient may sensitize them to chemotherapeutic agents. Indeed, some *RAD51* inhibitors have proven effective against certain cancers (27). In ccRCC Liu and Weng (28), 2022 stated that patients with low *RAD51* expression were found to be more likely to respond to immune checkpoint blockade immunotherapy. These findings indicate that *RAD51* protein may serve as a poor prognostic indicator and a potential target for developing anti-tumor strategies for ccRCC patients, necessitating further investigation.

BRCA1 and *BRCA2* are recognized as important tumor suppressor genes that help to prevent tumors by maintaining the stability of the genome. They perform complex tasks in the repair of DNA damage (29). *BRCA1* functions early in HR by detecting DNA damage and facilitating end resection, allowing ssDNA to bind replication protein A (RPA) before *RAD51* replaces RPA on the ssDNA. *BRCA2*, aided by *BRCA1* and *PALB2*, displaces RPA and loads *RAD51* onto the resected DNA, forming a *RAD51*-ssDNA filament that is essential for HR and for blocking the detrimental single-strand annealing pathway (30).

Although *BRCA1* and *BRCA2* mutations are mainly linked to ovarian and breast carcinomas (31,32), research on their role in ccRCC remains under active investigation. Understanding these associations could offer valuable insights for improved risk assessment, targeted therapies, and monitoring strategies for affected individuals.

In this study, high IHC expression of *BRCA1*, *BRCA2*, and co-expression of *BRCA1* and *BRCA2* in ccRCC patients were 51.3%, 35.5%, and 27.6%, respectively, and their expression was significantly associated with several favorable prognostic factors, including early tumor stage and absence of lymphatic

spread. Furthermore, survival analysis demonstrated an association between high *BRCA1* expression and longer OS and DFS in ccRCC patients. These findings indicate that *BRCA1* and *BRCA2* may play a role in suppressing progression of ccRCC and could be good prognostic indicators for patients with ccRCC.

In 2011, the first report documented a *BRCA1* germline mutation in a Pakistani patient with an aggressive form of ccRCC (33). In the Chinese population, inherited mutations in DDR-related genes, including *BRCA1*, have also been reported in RCC patients (34). A previous study reported a significant association between *BRCA1* alterations and the aggressiveness of ccRCC (35). In contrast, another study found no evidence that *BRCA1* mutations play a role in the development of kidney cancer in Polish individuals (36).

Recognizing VHL as a tumor suppressor gene in RCC (4), Scanlon et al. (37) reported that VHL-deficient cells in RCC had a decreased ability to carry out HR and reduced levels of specific HR genes, such as *BRCA1*. Additionally, VHL loss increased ER- α , which can bind to *BRCA1*, reduce *BRCA1* expression, block *BRCA1*-*RAD51* interaction, and induce γ -tubulin expression, which is strongly linked to resistance to microtubule-targeted therapy such as Taxol. This may represent a mechanism of drug resistance observed in both RCC and *BRCA1*-deficient cancers.

Diez-Calzadilla et al. (35) found that changes in *BRCA2* were linked to the development of RCC and were strongly associated with aggressive tumors, including higher Fuhrman grades and an increased risk of distant metastasis. Liu et al. (38) suggested that *BRCA2* may be linked to colorectal cancer, ovarian cancer, and RCC by interacting with downregulated BCCIP.

Souza's et al. (39) and Kong et al., (40) indicate that in certain RCCs with a particular morphology, a *BRCA2* mutation may serve as a key mutation. *BRCA2* mutations may contribute to tumor aggressiveness and progression to high-grade RCCs in RCC subtypes such as ccRCC and papillary RCC. Moreover, the chromosomal region containing *BRCA2* is often deleted in sarcomatoid RCC. These findings contradict the conclusions of Złowocka-Perłowska et al. (36), who stated that *BRCA2* does not appear to be associated with the development of kidney cancer. Because of the contradictory information on the involvement of *BRCA2* in RCC.

Mutations in *BRCA1/2* could increase the risk of developing RCC and may have contributed to genomic instability in ccRCC patients; further investigation is required to determine the impact of *BRCA1/2* mutations on ccRCC development.

From a therapeutic standpoint, tumors that harbor *BRCA1/2* germline mutations exhibit biological characteristics resulting in genomic instability, which may make them responsive to DNA-damaging therapies such as platinum-based chemotherapy

and poly (ADP-ribose) polymerase inhibitors (PARPi) (41). It will be interesting to investigate whether patients with ccRCC and *BRCA1/2* mutations could similarly benefit from these treatments.

In this research, we examined the role of *BRCA1* and *BRCA2* as cofactors for *RAD51* and found a significant indirect association between *RAD51* and *BRCA1* expression, suggesting the presence of unknown compensatory mechanisms that allow the maintenance or restoration of *RAD51* recruitment to DNA lesions even when either *BRCA1* or *BRCA2* is absent. Therefore, *RAD51* has a crucial role in the DNA repair mechanisms of most malignant tumors, regardless of whether they have normal or abnormal *BRCA* status (proficient or deficient) (19).

Study Limitations

Median follow-up is 27.4 months, which may be considered relatively short for metastatic ccRCC.

Conclusion

This study highlights the importance of HR proteins, namely *RAD51*, *BRCA1*, and *BRCA2*, in the development and progression of ccRCC. Increased *RAD51* expression and lower expression of *BRCA1* and *BRCA2* are linked to more aggressive tumors and shorter OS and DFS, indicating their potential as significant prognostic markers in IHC analyses and providing insights into novel treatment strategies for ccRCC patients.

Ethics

Ethics Committee Approval: The current study was permitted by the Ethical Committee of Zagazig University according to the Egyptian Ethical Guidelines (approval number: ZU-IRB#11335, date: 12.03.2024).

Informed Consent: Written consent was obtained from each member.

Footnotes

Authorship Contributions

Surgical and Medical Practices: A.M.A.E.M., H.M.A., E.G.A., A.H.H., H.L.M., Concept: A.M.A.E.M., Design: A.M.A.E.M., Data Collection or Processing: A.H.H., H.L.M., Analysis or Interpretation: H.M.A., Literature Search: H.M.A., Writing: A.M.A.E.M., H.M.A., H.L.M.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Siegel RL, Giaquinto AN, Jemal A. Cancer statistics, 2024. *CA Cancer J Clin.* 2024;74:12-49. Erratum in: *CA Cancer J Clin.* 2024;74:203. [\[Crossref\]](#)
2. Lobo J, Ohashi R, Amin MB, Berney DM, Comp  rat EM, Cree IA, Gill AJ, Hartmann A, Menon S, Netto GJ, Raspollini MR, Rubin MA, Tan PH, Tickoo SK, Tsuzuki T, Turajlic S, Zhou M, Srigley JR, Moch H. WHO 2022 landscape of papillary and chromophobe renal cell carcinoma. *Histopathology.* 2022;81:426-438. [\[Crossref\]](#)
3. Makhov P, Joshi S, Ghatalia P, Kutikov A, Uzzo RG, Kolenko VM. Resistance to systemic therapies in clear cell renal cell carcinoma: mechanisms and management strategies. *Mol Cancer Ther.* 2018;17:1355-1364. [\[Crossref\]](#)
4. de Cubas AA, Rathmell WK. Epigenetic modifiers: activities in renal cell carcinoma. *Nat Rev Urol.* 2018;15:599-614. [\[Crossref\]](#)
5. Ledermann JA, Drew Y, Kristeleit RS. Homologous recombination deficiency and ovarian cancer. *Eur J Cancer.* 2016;60:49-58. [\[Crossref\]](#)
6. P  dua JDB, Mariano CFA, Fabro AT, Tirapelli DPDC, Sankarankutty AK, Dos Santos JS, Brunaldi MO. Prognostic value of the immunohistochemical expression of RAD51 and BRCA2 in gastric adenocarcinoma. *J Histochem Cytochem.* 2022;70:199-210. [\[Crossref\]](#)
7. Gachechiladze M,   karda J, Soltermann A, Joerger M. RAD51 as a potential surrogate marker for DNA repair capacity in solid malignancies. *Int J Cancer.* 2017;141:1286-1294. [\[Crossref\]](#)
8. Li Y, Wang WY, Xiao JH, Xu F, Liao DY, Xie L, Wang J, Luo F. Overexpression of Rad51 predicts poor prognosis in colorectal cancer: our experience with 54 patients. *PLoS One.* 2017;12:e0167868. Retraction in: *PLoS One.* 2018;13:e0206398. [\[Crossref\]](#)
9. Qiao GB, Wu YL, Yang XN, Zhong WZ, Xie D, Guan XY, Fischer D, Kolberg HC, Kruger S, Stuerzbecher HW. High-level expression of Rad51 is an independent prognostic marker of survival in non-small-cell lung cancer patients. *Br J Cancer.* 2005;93:137-143. [\[Crossref\]](#)
10. Maacke H, Opitz S, Jost K, Hamdorf W, Henning W, Kr  ger S, Feller AC, Lopens A, Diedrich K, Schwinger E, St  rzbecher HW. Over-expression of wild-type Rad51 correlates with histological grading of invasive ductal breast cancer. *Int J Cancer.* 2000;88:907-913. [\[Crossref\]](#)
11. Inoue K, Fry EA. Haploinsufficient tumor suppressor genes. *Adv Med Biol.* 2017 1st Quarter;118:83-122. [\[Crossref\]](#)
12. Zhao W, Steinfeld JB, Liang F, Chen X, Maranon DG, Jian Ma C, Kwon Y, Rao T, Wang W, Sheng C, Song X, Deng Y, Jimenez-Sainz J, Lu L, Jensen RB, Xiong Y, Kupfer GM, Wiese C, Greene EC, Sung P. BRCA1-BARD1 promotes RAD51-mediated homologous DNA pairing. *Nature.* 2017 Oct 19;550(7676):360-365. [\[Crossref\]](#)
13. Kwon Y, R  sner H, Zhao W, Selemenakis P, He Z, Kawale AS, Katz JN, Rogers CM, Neal FE, Badamchi Shabestari A, Petrosius V, Singh AK, Joel MZ, Lu L, Holloway SP, Burma S, Mukherjee B, Hromas R, Mazin A, Wiese C, S  rensen CS, Sung P. DNA binding and RAD51 engagement by the BRCA2 C-terminus orchestrate DNA repair and replication fork preservation. *Nat Commun.* 2023;14:432. [\[Crossref\]](#)
14. Abul-Husn NS, Soper ER, Odogis JA, Cullina S, Bobo D, Moscati A, Rodriguez JE; CBIPM Genomics Team; Regeneron Genetics Center; Loos RJF, Cho JH, Belbin GM, Suckiel SA, Kenny EE. Exome sequencing reveals a high prevalence of BRCA1 and BRCA2 founder variants in a diverse population-based biobank. *Genome Med.* 2020;12:2. [\[Crossref\]](#)
15. Du C, Peng Y, He Y, Chen G, Chen H. Low levels of BRCA1 protein expression predict a worse prognosis in stage I-II colon cancer. *Int J Biol Markers.* 2021;36:47-53. [\[Crossref\]](#)
16. Jiang M, Jia K, Wang L, Li W, Chen B, Liu Y, Wang H, Zhao S, He Y, Zhou C. Alterations of DNA damage response pathway: biomarker and therapeutic strategy for cancer immunotherapy. *Acta Pharm Sin B.* 2021;11:2983-2994. [\[Crossref\]](#)

17. Matos-Rodrigues G, Guirouilh-Barbat J, Martini E, Lopez BS. Homologous recombination, cancer and the 'RAD51 paradox'. *NAR Cancer*. 2021;3:zcab016. [\[Crossref\]](#)
18. Lu H, Li Z, Liu L, Tao Y, Zhou Y, Mao X, Zhu A, Wu H, Zheng X. A pan-cancer analysis of the oncogenic roles of RAD51 in human tumors. *Advanced Gut & Microbiome Research*. 2022;1-28. [\[Crossref\]](#)
19. Orhan E, Velazquez C, Tabet I, Sardet C, Theillet C. Regulation of RAD51 at the transcriptional and functional levels: what prospects for cancer therapy? *Cancers (Basel)*. 2021;13:2930. [\[Crossref\]](#)
20. Alshareeda AT, Negm OH, Aleskandarany MA, Green AR, Nolan C, TigHhe PJ, Madhusudan S, Ellis IO, Rakha EA. Clinical and biological significance of RAD51 expression in breast cancer: a key DNA damage response protein. *Breast Cancer Res Treat*. 2016;159:41-53. [\[Crossref\]](#)
21. Gachechiladze M, Škarda J, Kolek V, Grygárková I, Langová K, Bouchal J, Kolář Z, Baty F, Stahel R, Weder W, Soltermann A, Joerger M. Prognostic and predictive value of loss of nuclear RAD51 immunoreactivity in resected non-small cell lung cancer patients. *Lung Cancer*. 2017;105:31-38. [\[Crossref\]](#)
22. Xu Y, Chen K, Cai Y, Cheng C, Zhang Z, Xu G. Overexpression of Rad51 predicts poor prognosis and silencing of Rad51 increases chemo-sensitivity to doxorubicin in neuroblastoma. *Am J Transl Res*. 2019;11:5788-5799. [\[Crossref\]](#)
23. Wu R, Patel A, Tokumaru Y, Asaoka M, Oshi M, Yan L, Ishikawa T, Takabe K. High RAD51 gene expression is associated with aggressive biology and with poor survival in breast cancer. *Breast Cancer Res Treat*. 2022;193:49-63. [\[Crossref\]](#)
24. Kim PM, Allen C, Wagener BM, Shen Z, Nickoloff JA. Overexpression of human RAD51 and RAD52 reduces double-strand break-induced homologous recombination in mammalian cells. *Nucleic Acids Res*. 2001;29:4352-4360. [\[Crossref\]](#)
25. Zhang X, Ma N, Yao W, Li S, Ren Z. RAD51 is a potential marker for prognosis and regulates cell proliferation in pancreatic cancer. *Cancer Cell Int*. 2019;19:356. [\[Crossref\]](#)
26. Wiegman AP, Al-Ejeh F, Chee N, Yap PY, Gorski JJ, Da Silva L, Bolderson E, Chenevix-Trench G, Anderson R, Simpson PT, Lakhani SR, Khanna KK. Rad51 supports triple negative breast cancer metastasis. *Oncotarget*. 2014;5:3261-3272. [\[Crossref\]](#)
27. Grundy MK, Buckanovich RJ, Bernstein KA. Regulation and pharmacological targeting of RAD51 in cancer. *NAR Cancer*. 2020;2:zca024. [\[Crossref\]](#)
28. Liu H, Weng J. A pan-cancer bioinformatic analysis of RAD51 regarding the values for diagnosis, prognosis, and therapeutic prediction. *Front Oncol*. 2022;12:858756. [\[Crossref\]](#)
29. Jeggo PA, Pearl LH, Carr AM. DNA repair, genome stability and cancer: a historical perspective. *Nat Rev Cancer*. 2016;16:35-42. [\[Crossref\]](#)
30. Yang H, Li Q, Fan J, Holloman WK, Pavletich NP. The BRCA2 homologue Brh2 nucleates RAD51 filament formation at a dsDNA-ssDNA junction. *Nature*. 2005;433:653-657. [\[Crossref\]](#)
31. Kechin AA, Boyarskikh UA, Ermolenko NA, Tyulyandina AS, Lazareva DG, Avdalyan AM, Tyulyandin SA, Kushlinskii NE, Filipenko ML. Loss of heterozygosity in BRCA1 and BRCA2 genes in patients with ovarian cancer and probability of its use for clinical classification of variations. *Bull Exp Biol Med*. 2018;165:94-100. [\[Crossref\]](#)
32. Semmler L, Reiter-Brennan C, Klein A. BRCA1 and breast cancer: a review of the underlying mechanisms resulting in the tissue-specific tumorigenesis in mutation carriers. *J Breast Cancer*. 2019;22:1-14. [\[Crossref\]](#)
33. Rashid MU, Gull S, Faisal S, Khaliq S, Asghar K, Siddiqui N, Amin A, Hamann U. Identification of the deleterious 2080insA BRCA1 mutation in a male renal cell carcinoma patient from a family with multiple cancer diagnoses from Pakistan. *Fam Cancer*. 2011;10:709-712. [\[Crossref\]](#)
34. Kong W, Yang T, Wen X, Mu Z, Zhao C, Han S, Tian J, Zhang X, Zhou T, Zhang Y, Lou F, Cao S, Wang H, Zhang J. Germline mutation landscape and associated clinical characteristics in Chinese patients with renal cell carcinoma. *Front Oncol*. 2021;11:737547. [\[Crossref\]](#)
35. Diez-Calzadilla NA, Noguera Salvá R, Soriano Sarrió P, Martínez-Jabaloyas JM. Genetic profile and immunohistochemical study of clear cell renal carcinoma: pathological-anatomical correlation and prognosis. *Cancer Treat Res Commun*. 2021;27:100374. [\[Crossref\]](#)
36. Złowocka-Perłowska E, Tołoczko-Grabarek A, Narod SA, Lubiński J. Germline BRCA1 and BRCA2 mutations and the risk of bladder or kidney cancer in Poland. *Hered Cancer Clin Pract*. 2022;20:13. [\[Crossref\]](#)
37. Scanlon SE, Hegan DC, Sulkowski PL, Glazer PM. Suppression of homology-dependent DNA double-strand break repair induces PARP inhibitor sensitivity in VHL -deficient human renal cell carcinoma. *Oncotarget*. 2018;9:4647-4660. [\[Crossref\]](#)
38. Liu X, Cao L, Ni J, Liu N, Zhao X, Wang Y, Zhu L, Wang L, Wang J, Yue Y, Cai Y, Jin J. Differential BCCIP gene expression in primary human ovarian cancer, renal cell carcinoma and colorectal cancer tissues. *Int J Oncol*. 2013;43:1925-1934. [\[Crossref\]](#)
39. Souza VC, de Freitas Vinhas C, Miguel D, Athanazio DA, Trpkov K. Renal cell carcinoma morphologically similar to fumarate hydratase-deficient RCC in a patient with BRCA2 germline mutation. *Pathol Int*. 2018;68:541-542. [\[Crossref\]](#)
40. Kong Q, Gagan J, Zhou M. A high-grade renal cell carcinoma with somatic BRCA2 mutation. *Pathol Int*. 2019;69:432-433. [\[Crossref\]](#)
41. Nientiedt C, Tolstov Y, Volckmar AL, Endris V, Bonekamp D, Haberkorn U, Jäger D, Sültmann H, Stenzinger A, Hohenfellner M, Grulich C, Duensing S. PARP inhibition in BRCA2-mutated prostate cancer. *Ann Oncol*. 2017;28:189-191. [\[Crossref\]](#)